



6 Foods That Could Be Triggering Your OAB Symptoms

by KRYSTINA OSTERMEYER

Foods to Avoid With Overactive Bladder

We've all heard this old adage, "You are what you eat." We know that food is medicine – it can be healing. Likewise, it can also be harmful to our bodies. If you have overactive bladder (OAB), the foods that you eat can be harmful. They can wreak harm on your genitourinary system. There are also certain foods to avoid with overactive bladder

Today, we will discuss foods that should be avoided if you have OAB, and we will also discuss foods that you should include in your diet that may be helpful.

8 Foods and Beverages to Avoid With OAB

There are a variety of triggers that are known to cause issues for OAB sufferers; keep in mind that these foods and beverages are common triggers – but they do not cause problems for everyone with OAB.

You may also find that there is some unusual food that triggers your OAB symptoms; figuring out what causes your OAB symptoms is key.

1. Acidic and Spicy Foods

One of the most common offenders is acidic foods. This includes foods like citrus fruits (lemons, limes and oranges), tomatoes and tomato products (such as salsa and pizza sauce).

Foods that are extremely spicy are also known culprits. Heavy spiced foods are included in this category as well, as foods such as chilies and wasabi.

2. Artificial Sweeteners

This includes aspartame and saccharin, as well as other artificial sweeteners.

Diet sodas are common offenders; be sure to watch for foods that are labeled as low fat, fat-free and diet, as they are often sweetened with artificial sweeteners. Watch your labels!

3. Gluten

If you also have a gluten sensitivity or have celiac disease, ensure that you are removing gluten from your diet. Consuming gluten when you have a gluten sensitivity or celiac disease can contribute to OAB symptoms.

If you have removed other common OAB triggers and are still having symptoms, you may want to consider discussing being tested for a gluten sensitivity. Gluten is found in barley, rye and wheat products.

4. Caffeinated Drinks

Coffee and tea should also be avoided. You may be thinking, “But how am I supposed to live without my daily coffee fix?” I hear you, friends – I hear you. But the caffeine content in coffee can exacerbate symptoms, such as increased urgency, urinary frequency and an increase in incontinence.

Choosing to reduce caffeine or switching to decaf may reduce your symptoms. You should keep in mind that even decaf coffee has a minimal amount of caffeine – that minimal amount may be enough to exacerbate symptoms in some people.

5. Chocolate

Since I just broke it to you that you may need to cut out – or at least cut back on – your coffee habit, I should tell you that chocolate also has caffeine. The caffeine content in chocolate may be enough to exacerbate symptoms in some OAB sufferers.

Dark chocolate has the most caffeine, so opt for white chocolate, which does not have any caffeine, or dark chocolate, which has more cocoa and can satisfy with a smaller portion.

6. Carbonated Beverages and Alcohol

Carbonated beverages can also wreak havoc on your bladder. This includes sodas, soda water, energy drinks and even champagne. The carbonation can be irritating to the bladder and champagne is alcohol, which is a known irritant to the bladder.

Alcohol can cause dehydration, which can irritate the bladder. Also, it can disrupt the signals to your brain that make you aware of bladder overflow.

Beverages and Foods That Are OAB-Friendly

If it seems as if your favorite foods are on the above list, you may need to reframe your thinking!

There are plenty of fantastic foods that you can include in your diet – many of which will not exacerbate your OAB symptoms and some of which are also rich in vitamins and minerals – an added bonus!

Non-acidic fruits include:

- Bananas.
- Grapes.
- Apples.
- Coconuts.
- Blackberries and strawberries.
- Watermelon.

Non-acidic vegetables include:

- Broccoli.
 - Asparagus.
 - Lettuce.
 - Peppers.
 - Kale.
 - Celery.
 - Carrots.
 - Cucumbers.
-

Foods that are high in fiber prevent constipation, and constipation can trigger OAB symptoms. Eating plenty of fiber-rich foods is important. The fruits and vegetables listed above are fiber-rich, as well as the foods listed below:

- Raspberries.
- Artichokes.
- Lentils.
- Beans.
- Oats.
- Barley.
- Bran.
- Almonds.

Fiber and Cranberry Juice

It is recommended to consume 20 to 26 grams of fiber a day or about 14 grams per 1,000 calories eaten.

Constipation can aggravate OAB and bladder incontinence by adding additional pressure against the bowel. If you have severe constipation, it can damage the nerves that support the pelvic floor muscles. If these muscles are not functionally, it can lead to bladder issues.

Ensuring your diet is fortified with the right amount of fiber will keep bowel problems in check and keep pelvic floor muscles healthy.

Now, moving onto cranberry juice. Cranberry juice is a type of juice, which is good for staving off urinary tract infection. It prevents bacteria from adhering to the lining of the bladder. It can be acidic, however, so you will need to determine if it aggravates your OAB.

The Importance of Hydration

It may seem counterintuitive, but drinking an adequate amount of water is important. You may be thinking, "That can't be right!"

A lot of people believe that restricting their fluid intake will prevent them from continually rushing to the bathroom. It sounds logical, right? This is false because when you drink too little, you become dehydrated.

When you become dehydrated, your urine becomes highly concentrated – this irritates the bladder, as well as increases the risk for a urinary tract infection.

Obviously, too much fluid intake wreaks havoc on your bladder too. An overfilled bladder will cause leakage of urine, as well as a continuous 'got to go' feeling.

Water is recommended. you do not necessarily need to drink eight glasses of water – just drink when you are thirsty. You can tell when you have had enough water based on the color of your urine – it should be clear or almost colorless in color.

The Bottom Line

There are plenty of changes you can make in your life if you want to manage your symptoms and it can start with the foods you ingest.

So, now that you know which foods to avoid with overactive bladder, it is time to take a good, hard look at the foods that you are eating. It may be helpful to keep a food log for several weeks in order to pinpoint a pattern.